

REFERENCES

- Allen, G.R. (1978) A review of the archerfishes (family Toxotidae). *Records of the Western Australian Museum*. **6**, 355-378.
- Allen, G.R. (1982) *Inland fishes of Western Australia*, p. 86. Western Australian Museum, Perth.
- Allen, G.R. (1989) *Freshwater fishes of Australia*, p. 240. T.F.H. Publications, New Jersey.
- Allen, G.R. (1991) *Field guide to the freshwater fishes of New Guinea*, p. 268. Christensen Research Institute, Madang.
- Allen, G.R. & Burgess, W.E. (1990) A review of the glassfishes (Chandidae) of Australia and New Guinea. *Records of the Western Australian Museum*. Supplement No. **34**, 139-206.
- Allen, G.R. & Cross, N.J. (1982) *Rainbowfishes of Australia and Papua New Guinea*, p. 141. T.F.H. Publications, New Jersey.
- Allen, G.R. & Feinberg, M.N. (1998) Descriptions of a new genus and four new species of freshwater catfishes (Plotosidae) from Australia. *Aqua, Journal of Ichthyology and Aquatic Biology*. **3**, 9-18.
- Allen, G.R. & Hoese, D.F. (1980) A collection of fishes from Cape York Peninsula, Australia. *Journal of the Royal Society of Western Australia*. **63**, 53-61.
- Allen, G.R. & Leggett, R. (1990) A collection of freshwater fishes from the Kimberley region of Western Australia. *Records of the Western Australian Museum*. **14**, 527-545.
- Allen, G.R. & Pusey, B.J. (in prep.) *Hephaestus tulliensis* De Vis, a valid species of grunter (Pisces: Terapontidae) from fresh waters of north-eastern Queensland, Australia.

- Atkins, B.P., Lloyd, L.N. & Gray, J.D. (1988) Fish. *Wetlands of the Bakers Range and Marcollat watercourses: environmental characteristics* (ed. by B. P. Atkins), pp. 89-95. Ecologic and Associates, for South Australian Department of Environment and Planning, Adelaide.
- Australian Water Resources Council. (1976) *Review of Australia's water resources 1975*, p. 170. Australian Government Publishing Service, Canberra.
- Banarescu, P. (1990) *Zoogeography of fresh waters. Volumes 1-3*, p. 1617. AULA-Verlag, Wiesbaden
- Bauer, A.F.T. (1955) *Atlas of Australian resources: drainage systems*. Division of National Mapping, Department of National Development, Canberra.
- Benbow, M.C. (1990) Tertiary coastal dunes of the Eucla Basin, Australia. *Geomorphology*. **3**, 9-29.
- Bennett, K.D. (1990) Milankovitch cycles and their effects on species in ecological and evolutionary time. *Paleobiology*. **16**, 11-21.
- Berra, T.M. & Weatherley, A.H. (1972) A systematic study of the Australian freshwater serranid fish genus *Maccullochella*. *Copeia*. **1972**, 53-64.
- Bierman, P. & Turner, J. (1995) ^{10}Be and ^{26}Al Evidence for exceptionally low rates of Australian bedrock erosion and the likely existence of pre-Pleistocene landscapes. *Quaternary Research*. **44**, 378-382.
- Bishop, P. (1982) Stability or change: a review of ideas on ancient drainage in eastern New South Wales. *Australian Geographer*. **15**, 219-230.
- Bishop, P. (1986) Horizontal stability of the Australian continental drainage divide in south central New South Wales during the Cainozoic. *Australian Journal of Earth Sciences*. **33**, 295-307.

- Bishop, P. (1988) The eastern highlands of Australia: the evolution of an intraplate highland belt. *Progress in Physical Geography*. **12**, 159-182.
- Bishop, P. (1995) Drainage rearrangement by river capture, beheading and diversion. *Progress in Physical Geography*. **19**, 449-473.
- Blake, D.H. & Ollier, C.D. (1969) Geomorphological evidence of Quaternary tectonics in Southwestern Papua. *Revue de Geomorphologie Dynamique*. **19**, 28-32.
- Bremer, K. (1994) Branch support and tree stability. *Cladistics*. **10**, 295-304.
- Brown, C.M. & Radke, B.M. (1989) Stratigraphy and sedimentology of Mid-Tertiary permeability barriers in the subsurface of the Murray Basin, southeastern Australia. *BMR Journal of Australian Geology and Geophysics*. **11**, 367-385.
- Brown, D.E., Reichenbacher, F. & Franson, S.E. (1998) *A classification of North American biotic communities*, p. 141. The University of Utah Press, Salt Lake City.
- Burckle, L.H. (1995) Current issues of Pliocene paleoclimatology. *Paleoclimate and evolution, with emphasis on human origins* (ed. by E. S. Vrba, G. H. Denton, T. C. Partridge and L. H. Burckle), pp. 3-7. Yale University Press, New Haven.
- Cadwallader, P.L. (1981) Past and present distributions and translocations of Macquarie perch *Macquaria australasica* (Pisces: Percichthyidae), with particular reference to Victoria. *Proceedings of the Royal Society of Victoria*. **93**, 23-30.
- Cadwallader, P.L. & Backhouse, G.N. (1983) *A guide to the freshwater fish of Victoria*, p. 249. Atkinson Government Printer, Melbourne.
- Chen, X.Y. & Barton, C.E. (1991) Onset of aridity and dune-building in central Australia: sedimentological and magnetostratigraphic evidence from Lake Amadeus. *Palaeogeography, Palaeoclimatology, Palaeoecology*. **84**, 55-73.

- Chenoweth, S.F. & Hughes, J.M. (1997) Genetic population structure of the catadromous Perciform: *Macquaria novemaculeata* (Percichthyidae). *Journal of Fish Biology*. **50**, 721-733.
- Chilcott, S.J. & Humphries, P. (1996) Freshwater fish of northeast Tasmania with notes on the dwarf galaxias. *Records of the Queen Victoria Museum and Art Gallery*. **103**, 145-149.
- Clarke, J.D.A. (1994) Evolution of the Lefroy and Cowan Palaeodrainage channels, Western Australia. *Australian Journal of Earth Sciences*. **41**, 55-68.
- Colhoun, E.A. & Fitzsimons, S.J. (1990) Late Cainozoic glaciation in western Tasmania, Australia. *Quaternary Science Reviews*. **9**, 199-216.
- Coventry, R.J., Stephenson, P.J. & Webb, A.W. (1985) Chronology of landscape evolution and soil development in the upper Flinders River area, Queensland, based on isotopic dating of Cainozoic basalts. *Australian Journal of Earth Sciences*. **32**, 433-447.
- Crowley, L.E.L.M. (1990) Biogeography of the endemic freshwater fish *Craterocephalus* (Family Atherinidae). *Memoirs of the Queensland Museum*. **28**, 89-98.
- Crowley, L.E.L.M. & Ivantsoff, W. (1988) A new species of Australian *Craterocephalus* (Pisces: Atherinidae) and redescription of four other species. *Records of the Western Australian Museum*. **14**, 151-169.
- Crowley, L.E.L.M. & Ivantsoff, W. (1990) A review of species previously identified as *Craterocephalus eyresii* (Pisces: Atherinidae). *Proceedings of the Linnean Society of New South Wales*. **112**, 87-103.
- Crowley, L.E.L.M., Ivantsoff, W. & Allen, G.R. (1986) Taxonomic position of two crimson-spotted rainbowfish, *Melanotaenia duboulayi* and *Melanotaenia fluviatilis* (Pisces: Melanotaeniidae), from eastern Australia, with special reference to their early life-history stages. *Australian Journal of Marine and Freshwater Research*. **37**, 385-398.

- Darlington, P.J. Jr. (1957) *Zoogeography: the geographic distribution of animals*, p. 675. J. Wiley & Sons, New York.
- Darlington, P.J. Jr. (1965) *Biogeography of the southern end of the world*, p. 236. Harvard University Press, Massachusetts.
- Doutch, H.F. (1972) The paleogeography of northern Australia and New Guinea and its relevance to the Torres Strait area. *Bridge and barrier: the natural and cultural history of Torres Strait* (ed. by D. Walker), pp. 1-10. Australian National University, Canberra.
- Dufty, S. (1986) Genetic and morphological divergence between populations of Macquarie perch (*Macquaria australasica*) east and west of the Great Dividing Range. Honours Thesis, University of New South Wales.
- Endler, J.A. (1982) Problems in distinguishing historical from ecological factors in biogeography. *American Zoology*. **22**, 441-452.
- Eschmeyer, W.M. (1998) *Catalog of fishes. Volumes 1-3*, p. 2905. Californian Academy of Sciences, San Francisco.
- Estes, R. (1984) Fish, amphibians and reptiles from the Etadunna Formation, Miocene of South Australia. *Australian Zoologist*. **21**, 335-343.
- Frakes, L.A., Burger, D., Apthorpe, M., Wiseman, J., Dettmann, M., Alley, N., Flint, R., Gravestock, D., Ludbrook, N., Backhouse, J., Skwarko, S., Scheibnerova, V., McMinn, A., Moore, P.S., Bolton, B.R., Douglas, J.G., Christ, R., Wade, M., Molnar, R.E., McGowran, B., Balme, B.E. & Day, R.A. (1987a) Australian Cretaceous shorelines, stage by stage. *Palaeogeography, Palaeoclimatology, Palaeoecology*. **59**, 31-48.
- Frakes, L.A., McGowran, B. & Bowler, J.M. (1987b) Evolution of Australian environments. *Fauna of Australia. Volume 1A* (ed. by G. R. Dyne and D. W. Walton), pp. 1-16. Australian Government Publishing Service, Canberra.

- Frankenberg, R. (1974) Native freshwater fish. *Monographiae Biologicae*. **25**, 113-140.
- Gale, S.J. (1992) Long-term landscape evolution in Australia. *Earth Surface Processes and Landforms*. **17**, 323-343.
- Glover, C.J.M. (1982) Adaptations of fishes in arid Australia. *Evolution of the flora and fauna of arid Australia* (ed. by W. R. Barker and P. J. M. Greenslade), pp. 241-246. Peacock Publications, Adelaide.
- Glover, C.J.M. (1989) Fishes. *Natural history of Dalhousie Springs* (ed. by W. Zeidler and W. F. Ponder), pp. 89-112. South Australian Museum, Adelaide.
- Glover, C.J.M. (1990) Fishes. *Natural history of the north-east deserts* (ed. by M. J. Tyler, C. R. Twidale, M. Davies and C. B. Wells), pp. 189-198. Royal Society of South Australia, Adelaide.
- Glover, C.J.M. & Sim, T.C. (1978) Studies on central Australian fishes: a progress report. *South Australian Naturalist*. **52**, 35-44.
- Grande, T. (1996) The interrelationships of fossil and Recent gonorynchid fishes with comments on two Cretaceous taxa from Israel. *Mesozoic fishes - systematics and paleoecology* (ed. by G. Arratia and G. Viohl), pp. 299-318. Verlag Dr. Friedrich Pfeil, München.
- Grimes, C.B. & Kingsford, M.J. (1996) How do riverine plumes of different sizes influence fish larvae: do they enhance recruitment? *Marine and Freshwater Research*. **47**, 191-208.
- Gudger, E.W. (1929) More rains of fishes. *Annals and Magazine of Natural History*. **(10)3**, 1-26.
- Hadfield, A.J., Ivantsoff, W. & Johnston, P.G. (1979) Clinal variation in electrophoretic and morphological characters between two nominal species of the genus

Pseudomugil (Pisces: Atheriniformes: Pseudomugilidae). *Australian Journal of Marine and Freshwater Research*. **30**, 375-386.

Hamlyn-Harris, R. (1930) The relative value of larval destructors and the part they play in mosquito control in Queensland. *Proceedings of the Royal Society of Queensland*. **41**, 23-38.

Haq, B.U., Hardenbol, J. & Vail, P.R. (1987) Chronology of fluctuating sea levels since the Triassic. *Science*. **235**, 1156-1167.

Haworth, R.J. & Ollier, C.D. (1992) Continental rifting and drainage reversal: the Clarence River of eastern Australia. *Earth Surface Processes and Landforms*. **17**, 387-397.

Henstridge, D.A. & Missen, D.D. (1982) Geology of oil-shale deposits with The Narrows graben, Queensland, Australia. *The American Association of Petroleum Geologists Bulletin*. **66**, 719-731.

Herbert, B.W., Peeters, J.A., Graham, P.A. & Hogan, A.E. (1995) *Freshwater fish and aquatic habitat survey of Cape York Peninsula*, p. 376. Cape York Peninsula Land Use Strategy, Office of the Co-ordinator General of Queensland, Brisbane, Department of the Environment, Sport and Territories, Canberra, Queensland Department of Primary Industries, Brisbane.

Hill, D., Playford, G. & Woods, J.T. (1970) *Cainozoic fossils of Queensland*, p. 36. Queensland Palaeontolgraphical Society, Brisbane.

Hills, S. (1934) Tertiary fresh water fishes from southern Queensland. *Memoirs of the Queensland Museum*. **10**, 157-173.

Hills, S. (1943) Tertiary fresh-water fishes and crocodilian remains from Gladstone and Duaranga, Queensland. *Memoirs of the Queensland Museum*. **12**, 96-100.

- Hills, S. (1946) Fossil Murray cod (*Maccullochella macquariensis*) from diatomaceous earths in New South Wales. *Records of the Australian Museum*. **21**, 380-382.
- Hills, S. (1956) A brief review of Australian fossil vertebrates. *Studies on fossil vertebrates* (ed. by T. S. Westoll), pp. 86-107. Athlone Press, London.
- Hortle, K.G. & Pearson, R.G. (1990) Fauna of the Annan River system, far north Queensland, with reference to the impact of tin mining. I. Fishes. *Australian Journal of Marine and Freshwater Research*. **41**, 677-694.
- similarities between river basins. *Environmental Biology of Fishes*. **39**, 365-380.
- Horton, D. (1973) The concept of zoogeographic subregions. *Systematic Zoology*. **22**, 191-195.
- Humphries, C.J. & Parenti, L.R. (1986) *Cladistic biogeography*, p. 98. Oxford University Press, Oxford.
- Hurwood, D.A. & Hughes, J.M. (1998) Phylogeography of the freshwater fish, *Mogurnda adspersa*, in streams of northeastern Queensland, Australia: evidence for altered drainage patterns. *Molecular Ecology*. **7**, 1507-1517.
- Hutchins, J.B. (1981) Freshwater fish fauna of the Mitchell Plateau area, Kimberley, Western Australia. *Biological survey of Mitchell Plateau and Admiralty Gulf, Kimberley, Western Australia* (no author), pp. 229-247. Western Australian Museum, Perth.
- Iredale, T. & Whitley, G.P. (1938) The fluvifaunulae of Australia. *South Australian Naturalist*. **18**, 64-68.
- Jackson, P.D. & Davies, J.N. (1983) Survey of the fish fauna in the Grampians regions, south-western Victoria. *Proceedings of the Royal Society of Victoria*. **95**, 39-51.

- Jennings, J.N. (1967) Some karst areas of Australia. *Landform studies from Australia and New Guinea* (ed. by J. N. Jennings and J. A. Mabbutt), pp. 256-292. Australian National University Press, Canberra.
- Jerry, D.R. (1997) Population genetic structure of the catadromous Australian bass from throughout its range. *Journal of Fish Biology*. **51**, 909-920.
- Jerry, D.R. & Baverstock, P.R. (1998) Consequences of a catadromous life-strategy for levels of mitochondrial DNA differentiation among populations of the Australian bass, *Macquaria novemaculeata*. *Molecular Ecology*. **7**, 1003-1013.
- Jerry, D.R. & Cairns, S.C. (1998) Morphological variation in the catadromous Australian bass, from seven geographically distinct riverine drainages. *Journal of Fish Biology*. **52**, 829-843.
- Jerry, D.R. & Woodland, D.J. (1997) Electrophoretic evidence for the presence of the undescribed 'Bellinger' catfish (*Tandanus* sp.) (Teleostei : Plotosidae) in four New South Wales mid-northern coastal rivers. *Marine and Freshwater Research*. **48**, 235-240.
- Johnson, R.W. (1989) *Intraplate volcanism in eastern Australia and New Zealand*, p. 408. Cambridge University Press, United Kingdom.
- Jones, M.R. & Torgersen, T. (1988) Late Quaternary evolution of Lake Carpentaria on the Australia-New Guinea continental shelf. *Australian Journal of Earth Sciences*. **35**, 313-324.
- Keast, J.A. (1959) The reptiles of Australia. *Monographiae Biologicae*. **8**, 115-135.
- Kemp, A. (1982a) Australian Mesozoic and Cenozoic lungfish. *The fossil vertebrate record of Australia* (ed. by P. V. Rich and E. M. Thompson), pp. 133-139. Monash University Publications Committee, Melbourne.

- Kemp, A. (1982b) *Neoceratodus djelleh*, a new ceratodont lungfish from Duaringa, Queensland. *Alcheringa*. **6**, 151-155.
- Kemp, A. (1992) New cranial remains of neoceratodonts (Osteichthyes: Dipnoi) from the Late Oligocene to Middle Miocene of northern Australia, with comments on generic characters from Cenozoic dipnoans. *Journal of Vertebrate Paleontology*. **12**, 284-293.
- Kemp, A. (1993) *Ceratodus diutinus*, a new ceratodont from Cretaceous and Late Oligocene-Miocene deposits in Australia. *The Paleontological Society*. **67**, 883-888.
- Kemp, A. (1997a) Four species of *Metaceratodus* (Osteichthyes: Dipnoi, Family Ceratodontidae) from Australian Mesozoic and Cenozoic deposits. *Journal of Vertebrate Paleontology*. **17**, 26-33.
- Kemp, A. (1997b) A revision of Australian Mesozoic and Cenozoic lungfish of the family Neoceratodontidae (Osteichthyes: Dipnoi), with a description of four new species. *Journal of Paleontology*. **71**, 713-733.
- Kemp, A. & Molnar, R.E. (1981) *Neoceratodus forsteri* from the Lower Cretaceous of New South Wales, Australia. *Journal of Paleontology*. **55**, 211-217.
- Kershaw, A.P., Martin, H.A. & McEwen Mason, J.R.C. (1994) The Neogene: a period of transition. *History of the Australian vegetation: Cretaceous to Recent* (ed. by R. S. Hill), pp. 299-327. University of Cambridge Press, Cambridge.
- Kennard, M. (1992) A progress report to the Queensland National Parks and Wildlife Service (unpublished report)
- Kruskal, J.B. (1964) Multidimensional scaling by optimizing goodness of fit to a nonmetric hypothesis. *Psychometrika*. **29**, 1-25.

- Kuiter, R.H., Humphries, P.A. & Arthington, A.H. (1996) Family Nannoperidae pygmy perches. *Freshwater fishes of south-eastern Australia* 2nd edn, (ed. by R. M. McDowall), pp. 168-175. Reed Books, Sydney.
- Lake, J.S. (1971) *Freshwater fishes and rivers of Australia*, p. 61. Thomas Nelson Australia Ltd, Melbourne.
- Larson, H.K. & Martin, K.C. (1990) *Freshwater fishes of the Northern Territory*, p 102. Northern Territory Museum of Arts and Sciences, Darwin.
- Lawver, L.A., Gahagan, L.M. & Coffin, M.F. (1992) The development of paleoseaways around Antarctica. *The Antarctic paleoenvironment: a perspective on global changes* (ed. by J. P. Kennett and D. A. Warnke), pp. 7-30. American Geophysical Union, Washington D. C.
- Legendre, L. & Legendre, P. (1983) *Numerical ecology*, p. 419. Elsevier Scientific Publishing Company, Amsterdam.
- Li, G. (1994) Systematic position of the Australian fossil osteoglossid fish †*Phareodus* (= *Phareoides*) *queenslandicus* Hills. *Memoirs of the Queensland Museum*. **37**, 287-300.
- Llewellyn, L.C. (1983) The distribution of fish in New South Wales. *Australian Society for Limnology Special Publication No. 7*, 1-70.
- Longman, H.A. (1929) Palaeontological notes. Specimens from a well at Brigalow. *Memoirs of the Queensland Museum*. **9**, 247-251.
- Martin, H.A. (1994) Australian Tertiary phytogeography: evidence from palynology. *History of the Australian vegetation: Cretaceous to Recent* (ed. by R. S. Hill), pp. 104-142. University of Cambridge Press, Cambridge.

- McCulloch, A.R. (1925a) Fishes of Australia. *The illustrated Australian encyclopedia. Volume 1 A to Lys* (ed. by A. W. Jose and H. J. Carter), pp. 465-467. Angus & Robertson Ltd, Sydney.
- McCulloch, A.R. (1925b) Raining fishes. *Australian Museum Magazine*. **2**, 217-218.
- McDowall, R.M. (1978) Generalized tracks and dispersal in biogeography. *Systematic Zoology*. **27**, 88-104.
- McDowall, R.M. (1981) The relationships of Australian freshwater fishes. *Monographiae Biologicae*. **41**, 1253-1273.
- McDowall, R.M. (1990) *New Zealand freshwater fishes*, 2nd edn, p. 553. Heinemann Reed, Auckland.
- McDowall, R.M. & Frankenberg, R.S. (1981) The galaxiid fishes of Australia. *Records of the Australian Museum*. **33**, 443-605.
- McGowran, B. (1989) The later Eocene transgressions in southern Australia. *Alcheringa*. **3**, 45-68.
- McKay, R.J. & Johnson, J.W. (1990) The freshwater and estuarine fishes. *The Brisbane River* (ed. by P. Davie, E. Stock and D. Low Choy), pp. 153-166. Australian Littoral Society and Queensland Museum, Brisbane.
- Merrick, J.R. & Schmida, G.E. (1984) *Australian freshwater fishes: biology and management*, p. 409 Griffith Press Ltd., Netley.
- Midgley, S.H., Midgley, M. & Rowland, S.J. (1991) Fishes of the Bulloo-Bancannia drainage division. *Memoirs of the Queensland Museum*. **30**, 505-508.
- Molnar, R.E. & Kurz, C. (1997) The distribution of Pleistocene vertebrates on the eastern Darling Downs, based on the Queensland Museum collections. *Proceedings of the Linnean Society of New South Wales*. **117**, 107-134.

- Mooi, R.D. & Gill, A.C (1995) Association of epaxial musculature with dorsal-fin pterygiophores in acanthomorph fishes, and its phylogenetic significance. *Bulletin Of The Natural History Museum Zoology Series*. **61**, 121-137.
- Morgan, D.L., Gill, H.S. & Potter, I.C. (1998) Distribution, identification and biology of freshwater fishes in south-western Australia. *Records of the Western Australian Museum, Supplement No.* **56**, 1-97.
- Musyl, M.K. & Keenan, C.P. (1992) Population genetics and zoogeography of Australian freshwater golden perch, *Macquaria ambigua* (Richardson 1845) (Teleostei: Percichthyidae), and electrophoretic identification of a new species from the Lake Eyre Basin. *Australian Journal of Marine and Freshwater Research*. **43**, 1585-1601.
- Musyl, M.K. & Keenan, C.P. (1996) Evidence for cryptic speciation in Australian freshwater eel-tailed catfish, *Tandanus tandanus* (Teleostei: Plotosidae). *Copeia*. **1996**, 526-534.
- Myers, G.S. (1938) Freshwater fishes and west Indian zoogeography. *Smithsonian Report*. **1937**, 339-364.
- Nix, H.A. (1982) Environmental determinants of biogeography and evolution in Terra Australis. *Evolution of the flora and fauna of arid Australia* (ed. by W.R. Barker and P.J.M. Greenslade), pp. 47-66. Peacock Publications, Adelaide.
- Nott, J.F. (1992) Long-term drainage evolution in the Shoalhaven catchment, southeast highlands, Australia. *Earth Surface Processes and Landforms*. **17**, 361-374.
- Nott, J.F. (1995) The antiquity of landscapes on the North Australian Craton and the implications for theories of long-term landscape evolution. *Journal of Geology*. **103**, 19-33.
- Nott, J.F. (1996) Long-term landscape evolution on Groote Eylandt, Northern Territory. *AGSO Journal of Australian Geology & Geophysics*. **16**, 303-7.

- Nott, J.F., Idnurm, M. & Young, R.W. (1991) Sedimentology, weathering, age and geomorphological significance of Tertiary sediments on the far south coast of New South Wales. *Australian Journal of Earth Sciences*. **38**, 357-373.
- O'Conner, D. (1897) Fish acclimatisation in Queensland. *Proceedings of the Royal Society of Queensland*. **12**, 108-110.
- Obregón-Barboza, H., Contreras-Balderas, S. & Lozano-Vilano, M.L. (1994) The fishes of northern and central Veracruz, Mexico. *Hydrobiologia*. **286**, 79-95.
- Ollier, C.D. (1978) Tectonics and geomorphology of the Eastern Highlands. *Landform evolution in Australasia* (ed. by J. L. Davies and M. A. J. Williams), pp. 5-47. Australian National University Press, Canberra.
- Ollier, C.D. (1995) Tectonics and landscape evolution in southeast Australia. *Geomorphology*. **12**, 37-44.
- Ollier, C.D., Armidale, G., Gaunt, F.M. & Jurkowski, I. (1988) The Kimberley Plateau, Western Australia: a Precambrian erosion surface. *Zeitschrift für Geomorphologie*. **32**, 239-46.
- Ollier, C.D. & Pain, C.F. (1994) Landscape evolution and tectonics in southeastern Australia. *BMR Journal of Australian Geology and Geophysics*. **15**, 335-345.
- Partridge, T.C., Bond, G.C., Hartnady, C.J.H., deMenocal, P.B. & Ruddiman, W.F. (1995) Climatic effects of Late Neogene tectonism and volcanism. *Paleoclimate and evolution, with emphasis on human origins* (ed. by E. S. Vrba, G. H. Denton, T. C. Partridge and L. H. Burckle), pp. 8-23. Yale University Press, New Haven.
- Platnick, N.I. & Nelson, G. (1978) A method of analysis for historical background. *Systematic Zoology*. **27**, 1-16.
- Pledge, N.S. (1984) A new Miocene vertebrate faunal assemblage from the Lake Eyre Basin: a preliminary report. *Australian Zoologist*. **21**, 345-355.

- Pollard, D.A. (1980) Family Apogonidae cardinalfishes and mouthbrooders. *Freshwater fishes of south-eastern Australia*, 1st edn, (ed. by R. M. McDowall), pp. 156-157. A.H. & A.W. Reed Pty. Ltd., Sydney.
- Pollard, D.A. (1996) Family Apogonidae cardinalfishes and mouthbrooders. *Freshwater fishes of south-eastern Australia*, 2nd edn, (ed. by R. M. McDowall), pp. 181-182. Reed Books, Sydney.
- Pollard, D.A., Davis, T.L.O. & Llewellyn, L.C. (1996) Family Plotosidae eel-tailed catfishes. *Freshwater fishes of south-eastern Australia*, 2nd edn, (ed. by R. M. McDowall), pp. 109-113. Reed Books, Sydney.
- Pusey, B.J., Arthington, A.H. & Read, M.G. (1998) Freshwater fishes of the Burdekin River, Australia: biogeography, history and spatial variation in community structure. *Environmental Biology of Fishes*. **53**, 303-318.
- Pusey, B.J. & Kennard, M.J. (1996) Species richness and geographical variation in assemblage structure of the freshwater fish fauna of the wet tropics region of northern Queensland. *Marine and Freshwater Research*. **47**, 563-573.
- Reshetnikov, Y.S & Shakirova, F.M. (1993) A zoogeographical analysis of the ichthyofauna of central Asia including a list of freshwater fishes. *Journal of Ichthyology*. **33**, 99-110.
- Roberts, T.R. (1978) An ichthyological survey of the Fly River in Papua New-Guinea with descriptions of new species. *Smithsonian Contributions to Zoology*. **281**, 1-70.
- Rohlf, F.J. (1972) An empirical comparison of three ordination techniques in numerical taxonomy. *Systematic Zoology*. **21**, 271-280.
- Rohlf, F.J. (1997) *NTSYSpc, version 2.02*. Exeter Software, Applied Biostatistics Inc.
- Rohlf, F.J. & Fisher, D.R. (1968) Tests for hierarchical structure in random data sets. *Systematic Zoology*. **17**, 407-412.

- Rosen, B.R. (1984) Reef coral biogeography and climate through the late Cainozoic: just islands in the sun or a critical pattern of islands. Fossils and climate (ed. by P. J. Brenchley). *Geological Journal Special Issue*. **11**, 201-62.
- Rosen, B.R. (1985) Long-term geographical controls on regional diversity. *The Open University Geological Society, Journal*. **6**, 25-30.
- Rowland, S.J. (1993) *Maccullochella ikei*, an endangered species of freshwater cod (Pisces: Percichthyidae) from the Clarence River system, NSW and *M. peelii mariensis*, a new subspecies from the Mary River system Qld. *Records of the Australian Museum*. **45**, 121-145.
- Saeed, B., Ivantsoff, W & Allen, G.R. (1989) Taxonomic revision of the family Pseudomugilidae (Order Atheriniformes). *Australian Journal of Marine and Freshwater Research*. **40**, 719-787.
- Shi, G.R. (1993) Multivariate data analysis in palaeoecology and palaeobiogeography - a review. *Palaeogeography, Palaeoclimatology, Palaeoecology*. **105**, 199-234.
- Shipway, B. (1947) Rains of fishes? *Western Australian Naturalist*. **1**, 47-48.
- Spry, M.J., Gibson, D.L. & Eggleton, R.A. (1999) Tertiary evolution in the coastal lowlands and the Clyde River palaeovalley in southeast New South Wales. *Australian Journal of Earth Sciences*. **46**, 173-180.
- Stephenson, A.E. & Brown, C.M. (1989) The ancient Murray River system. *BMR Journal of Australian Geology and Geophysics*. **11**, 387-95.
- Swofford, D.L. (1998) *PAUP: phylogenetic analysis using parsimony, version 4.01b*. Illinois Natural History Survey, Champaign.
- Taylor, G. (1911) Physiography of eastern Australia. *Commonwealth Bureau of Meteorology Bulletin*. **8**, 1-18.

- Taylor, G. (1994) Landscapes of Australia: their nature and evolution. *History of the Australian vegetation: Cretaceous to Recent* (ed. by R. S. Hill), pp. 60-79. Cambridge University Press, Cambridge.
- Tedford, R.H., Archer, M., Bartholomai, A., Plane, M., Pledge, N.S., Rich, T., Rich, P. & Wells, R.T. (1977) The discovery of Miocene vertebrates, Lake Frome area, South Australia. *BMR Journal of Australian Geology and Geophysics*. **2**, 53-57.
- Torgersen, T., Hutchinson, M.F., Searle, D.E. & Nix, H.A. (1983) General bathymetry of the Gulf of Carpentaria and the Quaternary physiography of Lake Carpentaria. *Palaeogeography, Palaeoclimatology, Palaeoecology*. **41**, 207-225.
- Torgersen, T., Luly, J., De Deckker, P., Jones, M.R., Searle, D.E., Chivas, A.R. & Ullman, W.J. (1988) Late Quaternary environments of the Carpentaria Basin, Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology*. **67**, 245-261.
- Tulip, J.R., Taylor, G. & Truswell, E.M. (1982) Palynology of Tertiary Lake Bunyan, Cooma, New South Wales. *BMR Journal of Australian Geology and Geophysics*. **7**, 255-68.
- Turner, S. (1981) Queensland Museum Report. *Bulletin of the Society of Vertebrate Paleontology*. **122**, 39-40.
- Turner, S. (1982) A catalogue of fossil fish in Queensland. *Memoirs of the Queensland Museum*. **20**, 599-611.
- Twidale, C.R. (1966) Late Cainozoic activity of the Selwyn Upwarp, Northwest Queensland. *Journal of the Geological Society of Australia*. **13**, 491-4.
- Twidale, C.R. (1997) The great age of some Australian landforms: examples of, and possible explanations for, landscape longevity. *Palaeosurfaces: recognition reconstruction and palaeoenvironmental interpretation* (ed. by M. Widdowson), pp. 13-23. Geological Society of London, London.

- Twidale, C.R. & Campbell, E.M. (1995) Pre-Quaternary landforms in the low latitude context: the example of Australia. *Geomorphology*. **12**, 17-35.
- Unmack, P.J. (1995) Desert fishes down under. *Proceedings of the Desert Fishes Council, 1994*. **26**, 71-95.
- Unmack, P.J. (in press) The genus *Hypseleotris* of southeastern Australia: its identification and breeding biology. *Fishes of Sahul*. **14**.
- Van der Beek, P.A., Braun, J. & Lambeck, K. (1999) Post-Palaeozoic uplift history of southeastern Australia revisited: results from a process based model of landscape evolution. *Australian Journal of Earth Sciences*. **46**, 157-172.
- Van de Graaff, W.J.E., Crowe, R.W.A., Bunting, J.A., Jackson, P. & Jackson, M.J. (1977) Relict Early Cainozoic drainages in arid Western Australia. *Zeitschrift fur Geomorphologie*. **21**, 379-400.
- Vari, R.P. (1978) The terapon perches (Percoidei, Teraponidae). A cladistic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History*. **159**, 175-340.
- Veevers, J.J. (1984) *Phanerozoic earth history of Australia*, p. 418. Clarendon Press, Oxford.
- Veevers, J.J. (1991) Mid-Cretaceous tectonic climax, Late Cretaceous recovery, and Cainozoic relaxation in the Australian region. *The Cainozoic in Australia: a reappraisal of the evidence* (ed. by M. A. J. Williams, P. De Deckker and A. P. Kershaw), pp. 1-14. Geological Society of Australia, Sydney.
- Veevers, J.J. & Eittreim, S.L. (1988) Reconstruction of Antarctica and Australia at breakup (95±5 Ma) and before rifting (160 Ma). *Australian Journal of Earth Sciences*. **35**, 355-362.

- Veevers, J.J., Stagg, H.M.J., Willcox, J.B. & Davies, H.L. (1990) Pattern of slow seafloor spreading (<4mm/year) from breakup (96 Ma) to A20 (44.5 Ma) off the southern margin of Australia. *BMR Journal of Australian Geology and Geophysics*. **11**, 499-507.
- Vickers-Rich, P. & Molnar, R.E. (1996) The foot of a bird from the Eocene Redbank Plains Formation of Queensland, Australia. *Alcheringa*. **20**, 21-29.
- Wager, R. (1993) *The distribution and conservation status of Queensland freshwater fishes*, p.62. Department of Primary Industries Information Series QI93001. Department of Primary Industries, Brisbane.
- Wallace, A.R. (1876) *The geographical distribution of animals*, 1962 reprint, p. 503. Hafner Publishing Company, New York.
- Warren, M.L. Jr., Burr, B.M., Etnier, D.A. & Starnes, W.C. (1991) Fishes of Kentucky and Tennessee: a hierarchical classification of drainages. *Journal of the Tennessee Academy of Science*. **66**, 135-140.
- Wasson, R.J. (1982) Landform development in Australia. *Evolution of the flora and fauna of arid Australia* (ed. by W. R. Barker and P. J. M. Greenslade), pp. 23-33. Peacock Publications, Frewville, South Australia.
- Watanabe, K. (1998). Parsimony analysis of the distribution pattern of Japanese primary freshwater fishes, and its application to the distribution of the bagrid catfishes. *Ichthyological Research*. **45**, 259-270.
- Waters, J.M., Lintermans, M. & White, R.W.G. (1994) Mitochondrial DNA variation suggests river capture as a source of variance in *Gadopsis bispinosus* (Pisces: Gadopsidae). *Journal of Fish Biology*. **44**, 549-551.
- Webster, P.J. & Stretten, N.A. (1972) Aspects of Late Quaternary climate in tropical Australasia. *Bridge and barrier: the natural and cultural history of Torres Strait* (ed. by D. Walker), pp. 39-60. Australian National University, Canberra.

- Whitley, G.P. (1943) The fishes of New Guinea. *The Australian Museum Magazine*. **8**, 141-143.
- Whitley, G.P. (1959) The freshwater fishes of Australia. *Monographiae Biologicae*. **8**, 136-149.
- Whitley, G.P. (1972) Rains of fishes in Australia. *Australian Natural History*. **17**, 154-159.
- Williams, D.L.G. (1980) Catalogue of Pleistocene vertebrate fossils and sites in South Australia. *Transactions of the Royal Society of South Australia*. **104**, 101-115.
- Williams, G.E. & Goode, A.D.T. (1978) Possible western outlet for an ancient Murray River in South Australia. *Search*. **9**, 443-447.
- Williams, M.A.J. (1984) Cenozoic evolution of arid Australia. *Arid Australia* (ed. by H. C. Cogger and E. E. Cameron), pp. 59-77. Australia Museum, Sydney.
- Williams, N.J. (1970) A comparison of the two species of the genus *Percalates* Ramsey and Ogilby (Percomorphi: Macquariidae) and their taxonomy. *New South Wales State Fisheries Bulletin*. **11**, 1-59.
- Williams, W.D. & Allen, G.R. (1987) Origins and adaptations of the fauna of inland waters. *Fauna of Australia. Volume 1A* (ed. by G. R. Dyne and D. W. Walton), pp. 184-201. Australian Government Publishing Service, Canberra.
- Wolanski, E. & Jones, M. (1981) Physical properties of Great Barrier Reef lagoon waters near Townsville. I. Effects of Burdekin River floods. *Australian Journal of Marine and Freshwater Research*. **32**, 305-319.
- Woodward, A.S. (1902) On *Atherstonia australis* and *Ctenolates avus*, two new species of fossil fishes from New South Wales. *Records of the Geological Survey of New South Wales*. **7**, 88-91.

Young, R.W. (1978) The study of landform evolution in the Sydney region: a review. *Australian Geographer*. **14**, 71-93.

Young, R.W. & McDougall, I. (1993) Long-term landscape evolution: Early Miocene and modern rivers in southern New South Wales, Australia. *The Journal of Geology*. **101**, 35-49.

Zhisheng, A., Bowler, J.M., Opdyke, N.D., Macumber, P.G. & Firman, J.B. (1986) Palaeomagnetic stratigraphy of Lake Bungunna: Plio-Pleistocene precursor of aridity in the Murray Basin, southeastern Australia. *Palaeogeography, Palaeoclimatology, Palaeoecology*. **54**, 219-239.